

# Cell Physiology of Exercise

BIOL4026

View Online



[1]

BAAR, K. 2006. Training for Endurance and Strength. *Medicine & Science in Sports & Exercise*. 38, 11 (Nov. 2006), 1939–1944.  
DOI:<https://doi.org/10.1249/01.mss.0000233799.62153.19>.

[2]

Baar, K. and Hardie, D.G. 2008. Small molecules can have big effects on endurance. *Nature Chemical Biology*. 4, 10 (Oct. 2008), 583–584.  
DOI:<https://doi.org/10.1038/nchembio1008-583>.

[3]

Barrès, R. et al. 2012. Acute Exercise Remodels Promoter Methylation in Human Skeletal Muscle. *Cell Metabolism*. 15, 3 (Mar. 2012), 405–411.  
DOI:<https://doi.org/10.1016/j.cmet.2012.01.001>.

[4]

Bogdanis, G.C. et al. 1995. Recovery of power output and muscle metabolites following 30 s of maximal sprint cycling in man. *The Journal of Physiology*. 482, 2 (Jan. 1995), 467–480.  
DOI:<https://doi.org/10.1113/jphysiol.1995.sp020533>.

[5]

Boluyt, M.O. et al. 2006. Changes in the rat heart proteome induced by exercise training: Increased abundance of heat shock protein hsp20. *PROTEOMICS*. 6, 10 (May 2006), 3154–3169. DOI:<https://doi.org/10.1002/pmic.200401356>.

[6]

BOOTH, F.W. et al. 1998. Molecular and cellular adaptation of muscle in response to physical training. *Acta Physiologica Scandinavica*. 162, 3 (Feb. 1998), 343–350. DOI:<https://doi.org/10.1046/j.1365-201X.1998.0326e.x>.

[7]

Burniston, J.G. 2009. Adaptation of the rat cardiac proteome in response to intensity-controlled endurance exercise. *PROTEOMICS*. 9, 1 (Jan. 2009), 106–115. DOI:<https://doi.org/10.1002/pmic.200800268>.

[8]

Burniston, J.G. 2008. Changes in the rat skeletal muscle proteome induced by moderate-intensity endurance exercise. *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics*. 1784, 7–8 (Jul. 2008), 1077–1086. DOI:<https://doi.org/10.1016/j.bbapap.2008.04.007>.

[9]

Burstein, B. and Nattel, S. 2008. Atrial Fibrosis: Mechanisms and Clinical Relevance in Atrial Fibrillation. *Journal of the American College of Cardiology*. 51, 8 (Feb. 2008), 802–809. DOI:<https://doi.org/10.1016/j.jacc.2007.09.064>.

[10]

Bye, A. et al. 2008. Aerobic capacity-dependent differences in cardiac gene expression. *Physiological Genomics*. 33, 1 (Mar. 2008), 100–109. DOI:<https://doi.org/10.1152/physiolgenomics.00269.2007>.

[11]

Bye, A. et al. 2008. Gene expression profiling of skeletal muscle in exercise-trained and sedentary rats with inborn high and low VO<sub>2</sub>. *Physiological Genomics*. 35, 3 (Nov. 2008), 213–221. DOI:<https://doi.org/10.1152/physiolgenomics.90282.2008>.

[12]

Carè, A. et al. 2007. MicroRNA-133 controls cardiac hypertrophy. *Nature Medicine*. 13, 5

(May 2007), 613–618. DOI:<https://doi.org/10.1038/nm1582>.

[13]

Casey, A. et al. 1996. Creatine ingestion favorably affects performance and muscle metabolism during maximal exercise in humans. *American Journal of Physiology-Endocrinology and Metabolism*. 271, 1 (Jul. 1996), E31–E37. DOI:<https://doi.org/10.1152/ajpendo.1996.271.1.E31>.

[14]

Chien, K.R. 2007. Molecular medicine: MicroRNAs and the tell-tale heart. *Nature*. 447, 7143 (May 2007), 389–390. DOI:<https://doi.org/10.1038/447389a>.

[15]

Creemers, E.E.J.M. et al. 2003. Deficiency of TIMP-1 exacerbates LV remodeling after myocardial infarction in mice. *American Journal of Physiology-Heart and Circulatory Physiology*. 284, 1 (Jan. 2003), H364–H371. DOI:<https://doi.org/10.1152/ajpheart.00511.2002>.

[16]

Daniels, A. et al. 2009. Connective tissue growth factor and cardiac fibrosis. *Acta Physiologica*. 195, 3 (Mar. 2009), 321–338. DOI:<https://doi.org/10.1111/j.1748-1716.2008.01936.x>.

[17]

Di Biase, V. and Franzini-Armstrong, C. 2005. Evolution of skeletal type e-c coupling. *The Journal of Cell Biology*. 171, 4 (Nov. 2005), 695–704. DOI:<https://doi.org/10.1083/jcb.200503077>.

[18]

Diffie, G.M. 2004. Adaptation of Cardiac Myocyte Contractile Properties to Exercise Training. *Exercise and Sport Sciences Reviews*. 32, 3 (Jul. 2004), 112–119. DOI:<https://doi.org/10.1097/00003677-200407000-00007>.

[19]

Eto, Y. et al. 2000. Calcineurin Is Activated in Rat Hearts With Physiological Left Ventricular Hypertrophy Induced by Voluntary Exercise Training. *Circulation*. 101, 18 (May 2000), 2134–2137. DOI:<https://doi.org/10.1161/01.CIR.101.18.2134>.

[20]

Fernandes, T. et al. 2015. Aerobic exercise training promotes physiological cardiac remodeling involving a set of microRNAs. *American Journal of Physiology-Heart and Circulatory Physiology*. 309, 4 (Aug. 2015), H543–H552. DOI:<https://doi.org/10.1152/ajpheart.00899.2014>.

[21]

Hambrecht, R. et al. 2003. Regular Physical Activity Improves Endothelial Function in Patients With Coronary Artery Disease by Increasing Phosphorylation of Endothelial Nitric Oxide Synthase. *Circulation*. 107, 25 (Jul. 2003), 3152–3158. DOI:<https://doi.org/10.1161/01.CIR.0000074229.93804.5C>.

[22]

Haram, P.M. et al. 1AD. Adaptation of endothelium to exercise training: Insights from experimental studies. 13, (1AD), 336–346.

[23]

Haram, P.M. et al. 2006. Time-course of endothelial adaptation following acute and regular exercise. *European Journal of Cardiovascular Prevention & Rehabilitation*. 13, 4 (Aug. 2006), 585–591. DOI:<https://doi.org/10.1097/01.hjr.0000198920.57685.76>.

[24]

Hawley, J.A. et al. 2014. Integrative Biology of Exercise. *Cell*. 159, 4 (Nov. 2014), 738–749. DOI:<https://doi.org/10.1016/j.cell.2014.10.029>.

[25]

Hill, M. et al. 2003. Muscle satellite (stem) cell activation during local tissue injury and repair. *Journal of Anatomy*. 203, 1 (Jul. 2003), 89–99.

DOI:<https://doi.org/10.1046/j.1469-7580.2003.00195.x>.

[26]

Hsu, C.-P. et al. 2008. Extracellular Matrix Remodeling Attenuated After Experimental Postinfarct Left Ventricular Aneurysm Repair. *The Annals of Thoracic Surgery*. 86, 4 (Oct. 2008), 1243–1249. DOI:<https://doi.org/10.1016/j.athoracsur.2008.06.043>.

[27]

Iemitsu, M. et al. 2006. Activation pattern of MAPK signaling in the hearts of trained and untrained rats following a single bout of exercise. *Journal of Applied Physiology*. 101, 1 (Jul. 2006), 151–163. DOI:<https://doi.org/10.1152/jappphysiol.00392.2005>.

[28]

Iemitsu, M. et al. 2005. Gene expression profiling of exercise-induced cardiac hypertrophy in rats. *Acta Physiologica Scandinavica*. 185, 4 (Nov. 2005), 259–270. DOI:<https://doi.org/10.1111/j.1365-201X.2005.01494.x>.

[29]

Jørgensen, S.B. et al. 2006. Role of AMPK in skeletal muscle metabolic regulation and adaptation in relation to exercise. *The Journal of Physiology*. 574, 1 (Jul. 2006), 17–31. DOI:<https://doi.org/10.1113/jphysiol.2006.109942>.

[30]

KEMI, O. et al. 2007. Exercise training restores aerobic capacity and energy transfer systems in heart failure treated with losartan. *Cardiovascular Research*. 76, 1 (Oct. 2007), 91–99. DOI:<https://doi.org/10.1016/j.cardiores.2007.06.008>.

[31]

KEMI, O. et al. 2005. Moderate vs. high exercise intensity: Differential effects on aerobic fitness, cardiomyocyte contractility, and endothelial function. *Cardiovascular Research*. 67, 1 (Jul. 2005), 161–172. DOI:<https://doi.org/10.1016/j.cardiores.2005.03.010>.

[32]

Kemi, O.J. et al. 2008. Activation or inactivation of cardiac Akt/mTOR signaling diverges physiological from pathological hypertrophy. *Journal of Cellular Physiology*. 214, 2 (Feb. 2008), 316–321. DOI:<https://doi.org/10.1002/jcp.21197>.

[33]

Kemi, O.J. et al. 2004. Aerobic Fitness Is Associated With Cardiomyocyte Contractile Capacity and Endothelial Function in Exercise Training and Detraining. *Circulation*. 109, 23 (Jun. 2004), 2897–2904. DOI:<https://doi.org/10.1161/01.CIR.0000129308.04757.72>.

[34]

Kemi, O.J. et al. 2007. Aerobic interval training enhances cardiomyocyte contractility and Ca<sup>2+</sup> cycling by phosphorylation of CaMKII and Thr-17 of phospholamban. *Journal of Molecular and Cellular Cardiology*. 43, 3 (Sep. 2007), 354–361. DOI:<https://doi.org/10.1016/j.yjmcc.2007.06.013>.

[35]

Kemi, O.J. and Wisløff, U. 2010. Mechanisms of exercise-induced improvements in the contractile apparatus of the mammalian myocardium. *Acta Physiologica*. 199, 4 (Aug. 2010), 425–439. DOI:<https://doi.org/10.1111/j.1748-1716.2010.02132.x>.

[36]

Kiens, B. and Richter, E.A. 1998. Utilization of skeletal muscle triacylglycerol during postexercise recovery in humans. *American Journal of Physiology-Endocrinology and Metabolism*. 275, 2 (Aug. 1998), E332–E337. DOI:<https://doi.org/10.1152/ajpendo.1998.275.2.E332>.

[37]

Kong, S.W. et al. 2005. Genetic expression profiles during physiological and pathological cardiac hypertrophy and heart failure in rats. *Physiological Genomics*. 21, 1 (Mar. 2005), 34–42. DOI:<https://doi.org/10.1152/physiolgenomics.00226.2004>.

[38]

KOVANEN, V. et al. 1980. Connective tissue of "fast" and "slow" skeletal muscle in rats...effects of endurance training. *Acta Physiologica Scandinavica*. 108, 2 (Feb. 1980), 173–180. DOI:<https://doi.org/10.1111/j.1748-1716.1980.tb06515.x>.

[39]

Linke, A. et al. 1AD. Effects of exercise training upon endothelial function in patients with cardiovascular disease. 13, (1AD), 424–432.

[40]

Lundby, C. et al. 2017. Biology of VO<sub>2</sub> max: looking under the physiology lamp. *Acta Physiologica*. 220, 2 (Jun. 2017), 218–228. DOI:<https://doi.org/10.1111/apha.12827>.

[41]

Maillet, M. et al. 2013. Molecular basis of physiological heart growth: fundamental concepts and new players. *Nature Reviews Molecular Cell Biology*. 14, 1 (Jan. 2013), 38–48. DOI:<https://doi.org/10.1038/nrm3495>.

[42]

Meeusen, R. et al. 2004. Hormonal responses in athletes: the use of a two bout exercise protocol to detect subtle differences in (over)training status. *European Journal of Applied Physiology*. 91, 2–3 (Mar. 2004), 140–146. DOI:<https://doi.org/10.1007/s00421-003-0940-1>.

[43]

Miyachi, M. et al. 1998. Effects of endurance training on the size and blood flow of the arterial conductance vessels in humans. *Acta Physiologica Scandinavica*. 163, 1 (May 1998), 13–16. DOI:<https://doi.org/10.1046/j.1365-201x.1998.0337f.x>.

[44]

MURPHY, G. and NAGASE, H. 2008. Progress in matrix metalloproteinase research. *Molecular Aspects of Medicine*. 29, 5 (Oct. 2008), 290–308. DOI:<https://doi.org/10.1016/j.mam.2008.05.002>.

[45]

Ramey, D.W. 1999. How to Read a Scientific Paper. AAEP PROCEEDINGS.

[46]

Reid, M.B. 2005. Response of the ubiquitin-proteasome pathway to changes in muscle activity. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 288, 6 (Jun. 2005), R1423–R1431.  
DOI:<https://doi.org/10.1152/ajpregu.00545.2004>.

[47]

Rowe, G.C. et al. 2014. Running Forward. *Circulation*. 129, 7 (Feb. 2014), 798–810.  
DOI:<https://doi.org/10.1161/CIRCULATIONAHA.113.001590>.

[48]

Spence, A.L. et al. 2013. A prospective randomized longitudinal study involving 6 months of endurance or resistance exercise. Conduit artery adaptation in humans. *The Journal of Physiology*. 591, 5 (Mar. 2013), 1265–1275.  
DOI:<https://doi.org/10.1113/jphysiol.2012.247387>.

[49]

Tsintzas, O.K. et al. 1996. Carbohydrate ingestion and single muscle fiber glycogen metabolism during prolonged running in men. *Journal of Applied Physiology*. 81, 2 (Aug. 1996), 801–809. DOI:<https://doi.org/10.1152/jappl.1996.81.2.801>.

[50]

Walter, G. et al. 1997. Noninvasive measurement of phosphocreatine recovery kinetics in single human muscles. *American Journal of Physiology-Cell Physiology*. 272, 2 (Feb. 1997), C525–C534. DOI:<https://doi.org/10.1152/ajpcell.1997.272.2.C525>.

[51]

Wilkins, B.J. et al. 2004. Calcineurin/NFAT Coupling Participates in Pathological, but not



Physiological, Cardiac Hypertrophy. *Circulation Research*. 94, 1 (Jan. 2004), 110–118.  
DOI:<https://doi.org/10.1161/01.RES.0000109415.17511.18>.

[52]

Williams, P.E. and Goldspink, G. Connective tissue changes in immobilised muscle. 138, 2, 343–350.

[53]

Wisløff, U. 2002. Aerobic exercise reduces cardiomyocyte hypertrophy and increases contractility, Ca<sup>2+</sup> sensitivity and SERCA-2 in rat after myocardial infarction. *Cardiovascular Research*. 54, 1 (Apr. 2002), 162–174.  
DOI:[https://doi.org/10.1016/S0008-6363\(01\)00565-X](https://doi.org/10.1016/S0008-6363(01)00565-X).

[54]

Wisløff, U. et al. 2007. Superior Cardiovascular Effect of Aerobic Interval Training Versus Moderate Continuous Training in Heart Failure Patients. *Circulation*. 115, 24 (Jun. 2007), 3086–3094. DOI:<https://doi.org/10.1161/CIRCULATIONAHA.106.675041>.