

# Cell Physiology of Exercise

BIOL4026

View Online



1

Lundby C, Montero D, Joyner M. Biology of VO<sub>2</sub> max: looking under the physiology lamp. *Acta Physiologica* 2017;**220**:218–28. doi:10.1111/apha.12827

2

Ramey DW. How to Read a Scientific Paper. AAEP PROCEEDINGS 1999.  
<https://pdfs.semanticscholar.org/104b/3127547393d6b94a8641100e9c297d653f56.pdf>

3

BAAR K. Training for Endurance and Strength. *Medicine & Science in Sports & Exercise* 2006;**38**:1939–44. doi:10.1249/01.mss.0000233799.62153.19

4

Baar K, Hardie DG. Small molecules can have big effects on endurance. *Nature Chemical Biology* 2008;**4**:583–4. doi:10.1038/nchembio1008-583

5

Barrès R, Yan J, Egan B, et al. Acute Exercise Remodels Promoter Methylation in Human Skeletal Muscle. *Cell Metabolism* 2012;**15**:405–11. doi:10.1016/j.cmet.2012.01.001

6

Carè A, Catalucci D, Felicetti F, et al. MicroRNA-133 controls cardiac hypertrophy. *Nature Medicine* 2007;**13**:613–8. doi:10.1038/nm1582

7

Chien KR. Molecular medicine: MicroRNAs and the tell-tale heart. *Nature* 2007;**447**:389–90. doi:10.1038/447389a

8

Eto Y, Yonekura K, Sonoda M, et al. Calcineurin Is Activated in Rat Hearts With Physiological Left Ventricular Hypertrophy Induced by Voluntary Exercise Training. *Circulation* 2000;**101**:2134–7. doi:10.1161/01.CIR.101.18.2134

9

Fernandes T, Baraúna VG, Negrão CE, et al. Aerobic exercise training promotes physiological cardiac remodeling involving a set of microRNAs. *American Journal of Physiology-Heart and Circulatory Physiology* 2015;**309**:H543–52. doi:10.1152/ajpheart.00899.2014

10

Iemitsu M, Maeda S, Jesmin S, et al. Activation pattern of MAPK signaling in the hearts of trained and untrained rats following a single bout of exercise. *Journal of Applied Physiology* 2006;**101**:151–63. doi:10.1152/jappphysiol.00392.2005

11

Maillet M, van Berlo JH, Molkenin JD. Molecular basis of physiological heart growth: fundamental concepts and new players. *Nature Reviews Molecular Cell Biology* 2013;**14**:38–48. doi:10.1038/nrm3495

12

Wilkins BJ, Dai Y-S, Bueno OF, et al. Calcineurin/NFAT Coupling Participates in Pathological, but not Physiological, Cardiac Hypertrophy. *Circulation Research* 2004;**94**:110–8. doi:10.1161/01.RES.0000109415.17511.18

13

Boluyt MO, Brevick JL, Rogers DS, et al. Changes in the rat heart proteome induced by exercise training: Increased abundance of heat shock protein hsp20. *PROTEOMICS* 2006;**6**:3154–69. doi:10.1002/pmic.200401356

14

Burniston JG. Changes in the rat skeletal muscle proteome induced by moderate-intensity endurance exercise. *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics* 2008;**1784**:1077–86. doi:10.1016/j.bbapap.2008.04.007

15

Burniston JG. Adaptation of the rat cardiac proteome in response to intensity-controlled endurance exercise. *PROTEOMICS* 2009;**9**:106–15. doi:10.1002/pmic.200800268

16

Bye A, Langaas M, Høydal MA, et al. Aerobic capacity-dependent differences in cardiac gene expression. *Physiological Genomics* 2008;**33**:100–9. doi:10.1152/physiolgenomics.00269.2007

17

Bye A, Høydal MA, Catalucci D, et al. Gene expression profiling of skeletal muscle in exercise-trained and sedentary rats with inborn high and low VO. *Physiological Genomics* 2008;**35**:213–21. doi:10.1152/physiolgenomics.90282.2008

18

Iemitsu M, Maeda S, Miyauchi T, et al. Gene expression profiling of exercise-induced cardiac hypertrophy in rats. *Acta Physiologica Scandinavica* 2005;**185**:259–70. doi:10.1111/j.1365-201X.2005.01494.x

19

Kong SW, Bodyak N, Yue P, et al. Genetic expression profiles during physiological and pathological cardiac hypertrophy and heart failure in rats. *Physiological Genomics* 2005;**21**:34–42. doi:10.1152/physiolgenomics.00226.2004

20

Diffie GM. Adaptation of Cardiac Myocyte Contractile Properties to Exercise Training. *Exercise and Sport Sciences Reviews* 2004;**32**:112–9. doi:10.1097/00003677-200407000-00007

21

Kemi OJ, Haram PM, Wisløff U, et al. Aerobic Fitness Is Associated With Cardiomyocyte Contractile Capacity and Endothelial Function in Exercise Training and Detraining. *Circulation* 2004;**109**:2897–904. doi:10.1161/01.CIR.0000129308.04757.72

22

KEMI O, HARAM P, LOENNECHEN J, et al. Moderate vs. high exercise intensity: Differential effects on aerobic fitness, cardiomyocyte contractility, and endothelial function. *Cardiovascular Research* 2005;**67**:161–72. doi:10.1016/j.cardiores.2005.03.010

23

Kemi OJ, Ellingsen Ø, Ceci M, et al. Aerobic interval training enhances cardiomyocyte contractility and Ca<sup>2+</sup> cycling by phosphorylation of CaMKII and Thr-17 of phospholamban. *Journal of Molecular and Cellular Cardiology* 2007;**43**:354–61. doi:10.1016/j.yjmcc.2007.06.013

24

Kemi OJ, Ceci M, Wisloff U, et al. Activation or inactivation of cardiac Akt/mTOR signaling diverges physiological from pathological hypertrophy. *Journal of Cellular Physiology* 2008;**214**:316–21. doi:10.1002/jcp.21197

25

Kemi OJ, Wisløff U. Mechanisms of exercise-induced improvements in the contractile apparatus of the mammalian myocardium. *Acta Physiologica* 2010;**199**:425–39. doi:10.1111/j.1748-1716.2010.02132.x

26

Hsu C-P, Huang C-Y, Wang J-S, et al. Extracellular Matrix Remodeling Attenuated After

Experimental Postinfarct Left Ventricular Aneurysm Repair. *The Annals of Thoracic Surgery* 2008;**86**:1243–9. doi:10.1016/j.athoracsur.2008.06.043

27

Burstein B, Nattel S. Atrial Fibrosis: Mechanisms and Clinical Relevance in Atrial Fibrillation. *Journal of the American College of Cardiology* 2008;**51**:802–9. doi:10.1016/j.jacc.2007.09.064

28

KOVANEN V, SUOMINEN H, HEIKKINEN E. Connective tissue of "fast" and "slow" skeletal muscle in rats...effects of endurance training. *Acta Physiologica Scandinavica* 1980;**108**:173–80. doi:10.1111/j.1748-1716.1980.tb06515.x

29

Daniels A, van Bilsen M, Goldschmeding R, et al. Connective tissue growth factor and cardiac fibrosis. *Acta Physiologica* 2009;**195**:321–38. doi:10.1111/j.1748-1716.2008.01936.x

30

Creemers EEJM, Davis JN, Parkhurst AM, et al. Deficiency of TIMP-1 exacerbates LV remodeling after myocardial infarction in mice. *American Journal of Physiology-Heart and Circulatory Physiology* 2003;**284**:H364–71. doi:10.1152/ajpheart.00511.2002

31

Williams PE, Goldspink G. Connective tissue changes in immobilised muscle. ;**138**:343–50. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1164074/>

32

MURPHY G, NAGASE H. Progress in matrix metalloproteinase research. *Molecular Aspects of Medicine* 2008;**29**:290–308. doi:10.1016/j.mam.2008.05.002

33

Di Biase V, Franzini-Armstrong C. Evolution of skeletal type e-c coupling. *The Journal of Cell Biology* 2005;**171**:695–704. doi:10.1083/jcb.200503077

34

Meeusen R, Piacentini MF, Busschaert B, et al. Hormonal responses in athletes: the use of a two bout exercise protocol to detect subtle differences in (over)training status. *European Journal of Applied Physiology* 2004;**91**:140–6. doi:10.1007/s00421-003-0940-1

35

BOOTH FW, TSENG BS, FLUCK M, et al. Molecular and cellular adaptation of muscle in response to physical training. *Acta Physiologica Scandinavica* 1998;**162**:343–50. doi:10.1046/j.1365-201X.1998.0326e.x

36

Hill M, Wernig A, Goldspink G. Muscle satellite (stem) cell activation during local tissue injury and repair. *Journal of Anatomy* 2003;**203**:89–99. doi:10.1046/j.1469-7580.2003.00195.x

37

Reid MB. Response of the ubiquitin-proteasome pathway to changes in muscle activity. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* 2005;**288**:R1423–31. doi:10.1152/ajpregu.00545.2004

38

Hambrecht R, Adams V, Erbs S, et al. Regular Physical Activity Improves Endothelial Function in Patients With Coronary Artery Disease by Increasing Phosphorylation of Endothelial Nitric Oxide Synthase. *Circulation* 2003;**107**:3152–8. doi:10.1161/01.CIR.0000074229.93804.5C

39

Haram PM, Adams V, Kemi OJ, et al. Time-course of endothelial adaptation following acute and regular exercise. *European Journal of Cardiovascular Prevention & Rehabilitation* 2006;**13**:585–91. doi:10.1097/01.hjr.0000198920.57685.76

40

Haram PM, Kemi OJ, Wisloff U. Adaptation of endothelium to exercise training: Insights from experimental studies. *1AD*; **13**:336–46. <https://www.bioscience.org/2008/v13/af/2683/fulltext.htm>

41

Linke A, Erbs S, Hambrecht R. Effects of exercise training upon endothelial function in patients with cardiovascular disease. *1AD*; **13**:424–32. <https://www.bioscience.org/2008/v13/af/2689/fulltext.htm>

42

Miyachi M, Iemitsu M, Okutsu M, et al. Effects of endurance training on the size and blood flow of the arterial conductance vessels in humans. *Acta Physiologica Scandinavica* 1998; **163**:13–6. doi:10.1046/j.1365-201x.1998.0337f.x

43

Spence AL, Carter HH, Naylor LH, et al. A prospective randomized longitudinal study involving 6 months of endurance or resistance exercise. Conduit artery adaptation in humans. *The Journal of Physiology* 2013; **591**:1265–75. doi:10.1113/jphysiol.2012.247387

44

Bogdanis GC, Nevill ME, Boobis LH, et al. Recovery of power output and muscle metabolites following 30 s of maximal sprint cycling in man. *The Journal of Physiology* 1995; **482**:467–80. doi:10.1113/jphysiol.1995.sp020533

45

Casey A, Constantin-Teodosiu D, Howell S, et al. Creatine ingestion favorably affects performance and muscle metabolism during maximal exercise in humans. *American Journal of Physiology-Endocrinology and Metabolism* 1996; **271**:E31–7. doi:10.1152/ajpendo.1996.271.1.E31

46

Jørgensen SB, Richter EA, Wojtaszewski JFP. Role of AMPK in skeletal muscle metabolic regulation and adaptation in relation to exercise. *The Journal of Physiology* 2006;**574**:17–31. doi:10.1113/jphysiol.2006.109942

47

Kiens B, Richter EA. Utilization of skeletal muscle triacylglycerol during postexercise recovery in humans. *American Journal of Physiology-Endocrinology and Metabolism* 1998;**275**:E332–7. doi:10.1152/ajpendo.1998.275.2.E332

48

Tsintzas OK, Williams C, Boobis L, et al. Carbohydrate ingestion and single muscle fiber glycogen metabolism during prolonged running in men. *Journal of Applied Physiology* 1996;**81**:801–9. doi:10.1152/jappl.1996.81.2.801

49

Walter G, Vandeborne K, McCully KK, et al. Noninvasive measurement of phosphocreatine recovery kinetics in single human muscles. *American Journal of Physiology-Cell Physiology* 1997;**272**:C525–34. doi:10.1152/ajpcell.1997.272.2.C525

50

KEMI O, HOYDAL M, HARAM P, et al. Exercise training restores aerobic capacity and energy transfer systems in heart failure treated with losartan. *Cardiovascular Research* 2007;**76**:91–9. doi:10.1016/j.cardiores.2007.06.008

51

Wisløff U. Aerobic exercise reduces cardiomyocyte hypertrophy and increases contractility, Ca<sup>2+</sup> sensitivity and SERCA-2 in rat after myocardial infarction. *Cardiovascular Research* 2002;**54**:162–74. doi:10.1016/S0008-6363(01)00565-X

52

Wisløff U, Støylen A, Loennechen JP, et al. Superior Cardiovascular Effect of Aerobic Interval Training Versus Moderate Continuous Training in Heart Failure Patients. *Circulation*



2007;**115**:3086–94. doi:10.1161/CIRCULATIONAHA.106.675041

53

Hawley JA, Hargreaves M, Joyner MJ, et al. Integrative Biology of Exercise. *Cell* 2014;**159**:738–49. doi:10.1016/j.cell.2014.10.029

54

Rowe GC, Safdar A, Arany Z. Running Forward. *Circulation* 2014;**129**:798–810. doi:10.1161/CIRCULATIONAHA.113.001590