

Cell Physiology of Exercise

BIOL4026

View Online



[1]

C. Lundby, D. Montero, and M. Joyner, 'Biology of VO₂ max: looking under the physiology lamp', *Acta Physiologica*, vol. 220, no. 2, pp. 218–228, Jun. 2017, doi: 10.1111/apha.12827.

[2]

D. W. Ramey, How to Read a Scientific Paper, vol. 45. AAEP PROCEEDINGS, 1999, pp. 280–284. [Online]. Available: <https://pdfs.semanticscholar.org/104b/3127547393d6b94a8641100e9c297d653f56.pdf>

[3]

K. BAAR, 'Training for Endurance and Strength', *Medicine & Science in Sports & Exercise*, vol. 38, no. 11, pp. 1939–1944, Nov. 2006, doi: 10.1249/01.mss.0000233799.62153.19.

[4]

K. Baar and D. G. Hardie, 'Small molecules can have big effects on endurance', *Nature Chemical Biology*, vol. 4, no. 10, pp. 583–584, Oct. 2008, doi: 10.1038/nchembio1008-583.

[5]

R. Barrès et al., 'Acute Exercise Remodels Promoter Methylation in Human Skeletal Muscle', *Cell Metabolism*, vol. 15, no. 3, pp. 405–411, Mar. 2012, doi: 10.1016/j.cmet.2012.01.001.

[6]

A. Carè et al., 'MicroRNA-133 controls cardiac hypertrophy', *Nature Medicine*, vol. 13, no. 5, pp. 613–618, May 2007, doi: 10.1038/nm1582.

[7]

K. R. Chien, 'Molecular medicine: MicroRNAs and the tell-tale heart', *Nature*, vol. 447, no. 7143, pp. 389–390, May 2007, doi: 10.1038/447389a.

[8]

Y. Eto et al., 'Calcineurin Is Activated in Rat Hearts With Physiological Left Ventricular Hypertrophy Induced by Voluntary Exercise Training', *Circulation*, vol. 101, no. 18, pp. 2134–2137, May 2000, doi: 10.1161/01.CIR.101.18.2134.

[9]

T. Fernandes, V. G. Baraúna, C. E. Negrão, M. I. Phillips, and E. M. Oliveira, 'Aerobic exercise training promotes physiological cardiac remodeling involving a set of microRNAs', *American Journal of Physiology-Heart and Circulatory Physiology*, vol. 309, no. 4, pp. H543–H552, Aug. 2015, doi: 10.1152/ajpheart.00899.2014.

[10]

M. Iemitsu, S. Maeda, S. Jesmin, T. Otsuki, Y. Kasuya, and T. Miyauchi, 'Activation pattern of MAPK signaling in the hearts of trained and untrained rats following a single bout of exercise', *Journal of Applied Physiology*, vol. 101, no. 1, pp. 151–163, Jul. 2006, doi: 10.1152/japplphysiol.00392.2005.

[11]

M. Maillet, J. H. van Berlo, and J. D. Molkentin, 'Molecular basis of physiological heart growth: fundamental concepts and new players', *Nature Reviews Molecular Cell Biology*, vol. 14, no. 1, pp. 38–48, Jan. 2013, doi: 10.1038/nrm3495.

[12]

B. J. Wilkins et al., 'Calcineurin/NFAT Coupling Participates in Pathological, but not Physiological, Cardiac Hypertrophy', *Circulation Research*, vol. 94, no. 1, pp. 110–118, Jan.

2004, doi: 10.1161/01.RES.0000109415.17511.18.

[13]

M. O. Boluyt, J. L. Brevick, D. S. Rogers, M. J. Randall, A. F. Scalia, and Z. B. Li, 'Changes in the rat heart proteome induced by exercise training: Increased abundance of heat shock protein hsp20', *PROTEOMICS*, vol. 6, no. 10, pp. 3154–3169, May 2006, doi: 10.1002/pmic.200401356.

[14]

J. G. Burniston, 'Changes in the rat skeletal muscle proteome induced by moderate-intensity endurance exercise', *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics*, vol. 1784, no. 7–8, pp. 1077–1086, Jul. 2008, doi: 10.1016/j.bbapap.2008.04.007.

[15]

J. G. Burniston, 'Adaptation of the rat cardiac proteome in response to intensity-controlled endurance exercise', *PROTEOMICS*, vol. 9, no. 1, pp. 106–115, Jan. 2009, doi: 10.1002/pmic.200800268.

[16]

A. Bye et al., 'Aerobic capacity-dependent differences in cardiac gene expression', *Physiological Genomics*, vol. 33, no. 1, pp. 100–109, Mar. 2008, doi: 10.1152/physiolgenomics.00269.2007.

[17]

A. Bye et al., 'Gene expression profiling of skeletal muscle in exercise-trained and sedentary rats with inborn high and low VO', *Physiological Genomics*, vol. 35, no. 3, pp. 213–221, Nov. 2008, doi: 10.1152/physiolgenomics.90282.2008.

[18]

M. Iemitsu, S. Maeda, T. Miyauchi, M. Matsuda, and H. Tanaka, 'Gene expression profiling of exercise-induced cardiac hypertrophy in rats', *Acta Physiologica Scandinavica*, vol. 185, no. 4, pp. 259–270, Nov. 2005, doi: 10.1111/j.1365-201X.2005.01494.x.

[19]

S. W. Kong et al., 'Genetic expression profiles during physiological and pathological cardiac hypertrophy and heart failure in rats', *Physiological Genomics*, vol. 21, no. 1, pp. 34–42, Mar. 2005, doi: 10.1152/physiolgenomics.00226.2004.

[20]

G. M. Diffie, 'Adaptation of Cardiac Myocyte Contractile Properties to Exercise Training', *Exercise and Sport Sciences Reviews*, vol. 32, no. 3, pp. 112–119, Jul. 2004, doi: 10.1097/00003677-200407000-00007.

[21]

O. J. Kemi, P. M. Haram, U. Wisløff, and Ø. Ellingsen, 'Aerobic Fitness Is Associated With Cardiomyocyte Contractile Capacity and Endothelial Function in Exercise Training and Detraining', *Circulation*, vol. 109, no. 23, pp. 2897–2904, Jun. 2004, doi: 10.1161/01.CIR.0000129308.04757.72.

[22]

O. KEMI et al., 'Moderate vs. high exercise intensity: Differential effects on aerobic fitness, cardiomyocyte contractility, and endothelial function', *Cardiovascular Research*, vol. 67, no. 1, pp. 161–172, Jul. 2005, doi: 10.1016/j.cardiores.2005.03.010.

[23]

O. J. Kemi et al., 'Aerobic interval training enhances cardiomyocyte contractility and Ca²⁺ cycling by phosphorylation of CaMKII and Thr-17 of phospholamban', *Journal of Molecular and Cellular Cardiology*, vol. 43, no. 3, pp. 354–361, Sep. 2007, doi: 10.1016/j.yjmcc.2007.06.013.

[24]

O. J. Kemi et al., 'Activation or inactivation of cardiac Akt/mTOR signaling diverges physiological from pathological hypertrophy', *Journal of Cellular Physiology*, vol. 214, no. 2, pp. 316–321, Feb. 2008, doi: 10.1002/jcp.21197.

[25]

O. J. Kemi and U. Wisløff, 'Mechanisms of exercise-induced improvements in the contractile apparatus of the mammalian myocardium', *Acta Physiologica*, vol. 199, no. 4, pp. 425–439, Aug. 2010, doi: 10.1111/j.1748-1716.2010.02132.x.

[26]

C.-P. Hsu, C.-Y. Huang, J.-S. Wang, P.-C. Sun, and C.-C. Shih, 'Extracellular Matrix Remodeling Attenuated After Experimental Postinfarct Left Ventricular Aneurysm Repair', *The Annals of Thoracic Surgery*, vol. 86, no. 4, pp. 1243–1249, Oct. 2008, doi: 10.1016/j.athoracsur.2008.06.043.

[27]

B. Burstein and S. Nattel, 'Atrial Fibrosis: Mechanisms and Clinical Relevance in Atrial Fibrillation', *Journal of the American College of Cardiology*, vol. 51, no. 8, pp. 802–809, Feb. 2008, doi: 10.1016/j.jacc.2007.09.064.

[28]

V. KOVANEN, H. SUOMINEN, and E. HEIKKINEN, 'Connective tissue of "fast" and "slow" skeletal muscle in rats...effects of endurance training', *Acta Physiologica Scandinavica*, vol. 108, no. 2, pp. 173–180, Feb. 1980, doi: 10.1111/j.1748-1716.1980.tb06515.x.

[29]

A. Daniels, M. van Bilsen, R. Goldschmeding, G. J. van der Vusse, and F. A. van Nieuwenhoven, 'Connective tissue growth factor and cardiac fibrosis', *Acta Physiologica*, vol. 195, no. 3, pp. 321–338, Mar. 2009, doi: 10.1111/j.1748-1716.2008.01936.x.

[30]

E. E. J. M. Creemers et al., 'Deficiency of TIMP-1 exacerbates LV remodeling after myocardial infarction in mice', *American Journal of Physiology-Heart and Circulatory Physiology*, vol. 284, no. 1, pp. H364–H371, Jan. 2003, doi: 10.1152/ajpheart.00511.2002.

[31]

P. E. Williams and G. Goldspink, 'Connective tissue changes in immobilised muscle', vol. 138, no. 2, pp. 343–350, [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1164074/>

[32]

G. MURPHY and H. NAGASE, 'Progress in matrix metalloproteinase research', *Molecular Aspects of Medicine*, vol. 29, no. 5, pp. 290–308, Oct. 2008, doi: 10.1016/j.mam.2008.05.002.

[33]

V. Di Biase and C. Franzini-Armstrong, 'Evolution of skeletal type e-c coupling', *The Journal of Cell Biology*, vol. 171, no. 4, pp. 695–704, Nov. 2005, doi: 10.1083/jcb.200503077.

[34]

R. Meeusen, M. F. Piacentini, B. Busschaert, L. Buyse, G. De Schutter, and J. Stray-Gundersen, 'Hormonal responses in athletes: the use of a two bout exercise protocol to detect subtle differences in (over)training status', *European Journal of Applied Physiology*, vol. 91, no. 2–3, pp. 140–146, Mar. 2004, doi: 10.1007/s00421-003-0940-1.

[35]

F. W. BOOTH, B. S. TSENG, M. FLUCK, and J. A. CARSON, 'Molecular and cellular adaptation of muscle in response to physical training', *Acta Physiologica Scandinavica*, vol. 162, no. 3, pp. 343–350, Feb. 1998, doi: 10.1046/j.1365-201X.1998.0326e.x.

[36]

M. Hill, A. Wernig, and G. Goldspink, 'Muscle satellite (stem) cell activation during local tissue injury and repair', *Journal of Anatomy*, vol. 203, no. 1, pp. 89–99, Jul. 2003, doi: 10.1046/j.1469-7580.2003.00195.x.

[37]

M. B. Reid, 'Response of the ubiquitin-proteasome pathway to changes in muscle activity', *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, vol. 288, no. 6, pp. R1423–R1431, Jun. 2005, doi: 10.1152/ajpregu.00545.2004.

[38]

R. Hambrecht et al., 'Regular Physical Activity Improves Endothelial Function in Patients With Coronary Artery Disease by Increasing Phosphorylation of Endothelial Nitric Oxide Synthase', *Circulation*, vol. 107, no. 25, pp. 3152–3158, Jul. 2003, doi: 10.1161/01.CIR.0000074229.93804.5C.

[39]

P. M. Haram et al., 'Time-course of endothelial adaptation following acute and regular exercise', *European Journal of Cardiovascular Prevention & Rehabilitation*, vol. 13, no. 4, pp. 585–591, Aug. 2006, doi: 10.1097/01.hjr.0000198920.57685.76.

[40]

P. M. Haram, O. J. Kemi, and U. Wisloff, 'Adaptation of endothelium to exercise training: Insights from experimental studies', vol. 13, pp. 336–346, 1 AD, [Online]. Available: <https://www.bioscience.org/2008/v13/af/2683/fulltext.htm>

[41]

A. Linke, S. Erbs, and R. Hambrecht, 'Effects of exercise training upon endothelial function in patients with cardiovascular disease', vol. 13, pp. 424–432, 1 AD, [Online]. Available: <https://www.bioscience.org/2008/v13/af/2689/fulltext.htm>

[42]

M. Miyachi, M. Iemitsu, M. Okutsu, and S. Onodera, 'Effects of endurance training on the size and blood flow of the arterial conductance vessels in humans', *Acta Physiologica Scandinavica*, vol. 163, no. 1, pp. 13–16, May 1998, doi: 10.1046/j.1365-201x.1998.0337f.x.

[43]

A. L. Spence, H. H. Carter, L. H. Naylor, and D. J. Green, 'A prospective randomized longitudinal study involving 6 months of endurance or resistance exercise. Conduit artery adaptation in humans', *The Journal of Physiology*, vol. 591, no. 5, pp. 1265–1275, Mar. 2013, doi: 10.1113/jphysiol.2012.247387.

[44]

G. C. Bogdanis, M. E. Nevill, L. H. Boobis, H. K. Lakomy, and A. M. Nevill, 'Recovery of power output and muscle metabolites following 30 s of maximal sprint cycling in man.', *The Journal of Physiology*, vol. 482, no. 2, pp. 467–480, Jan. 1995, doi: 10.1113/jphysiol.1995.sp020533.

[45]

A. Casey, D. Constantin-Teodosiu, S. Howell, E. Hultman, and P. L. Greenhaff, 'Creatine ingestion favorably affects performance and muscle metabolism during maximal exercise in humans', *American Journal of Physiology-Endocrinology and Metabolism*, vol. 271, no. 1, pp. E31–E37, Jul. 1996, doi: 10.1152/ajpendo.1996.271.1.E31.

[46]

S. B. Jørgensen, E. A. Richter, and J. F. P. Wojtaszewski, 'Role of AMPK in skeletal muscle metabolic regulation and adaptation in relation to exercise', *The Journal of Physiology*, vol. 574, no. 1, pp. 17–31, Jul. 2006, doi: 10.1113/jphysiol.2006.109942.

[47]

B. Kiens and E. A. Richter, 'Utilization of skeletal muscle triacylglycerol during postexercise recovery in humans', *American Journal of Physiology-Endocrinology and Metabolism*, vol. 275, no. 2, pp. E332–E337, Aug. 1998, doi: 10.1152/ajpendo.1998.275.2.E332.

[48]

O. K. Tsintzas, C. Williams, L. Boobis, and P. Greenhaff, 'Carbohydrate ingestion and single muscle fiber glycogen metabolism during prolonged running in men', *Journal of Applied Physiology*, vol. 81, no. 2, pp. 801–809, Aug. 1996, doi: 10.1152/jappl.1996.81.2.801.

[49]

G. Walter, K. Vandenborne, K. K. McCully, and J. S. Leigh, 'Noninvasive measurement of phosphocreatine recovery kinetics in single human muscles', *American Journal of Physiology-Cell Physiology*, vol. 272, no. 2, pp. C525–C534, Feb. 1997, doi: 10.1152/ajpcell.1997.272.2.C525.

[50]

O. KEMI et al., 'Exercise training restores aerobic capacity and energy transfer systems in

heart failure treated with losartan', *Cardiovascular Research*, vol. 76, no. 1, pp. 91–99, Oct. 2007, doi: 10.1016/j.cardiores.2007.06.008.

[51]

U. Wisløff, 'Aerobic exercise reduces cardiomyocyte hypertrophy and increases contractility, Ca²⁺ sensitivity and SERCA-2 in rat after myocardial infarction', *Cardiovascular Research*, vol. 54, no. 1, pp. 162–174, Apr. 2002, doi: 10.1016/S0008-6363(01)00565-X.

[52]

U. Wisløff et al., 'Superior Cardiovascular Effect of Aerobic Interval Training Versus Moderate Continuous Training in Heart Failure Patients', *Circulation*, vol. 115, no. 24, pp. 3086–3094, Jun. 2007, doi: 10.1161/CIRCULATIONAHA.106.675041.

[53]

J. A. Hawley, M. Hargreaves, M. J. Joyner, and J. R. Zierath, 'Integrative Biology of Exercise', *Cell*, vol. 159, no. 4, pp. 738–749, Nov. 2014, doi: 10.1016/j.cell.2014.10.029.

[54]

G. C. Rowe, A. Safdar, and Z. Arany, 'Running Forward', *Circulation*, vol. 129, no. 7, pp. 798–810, Feb. 2014, doi: 10.1161/CIRCULATIONAHA.113.001590.