

Cell Physiology of Exercise

BIOL4026

View Online



-
1.
Lundby C, Montero D, Joyner M. Biology of VO₂ max: looking under the physiology lamp. *Acta Physiologica*. 2017 Jun;220(2):218–28.

 2.
Ramey DW. How to Read a Scientific Paper [Internet]. Vol. 45. AAEP PROCEEDINGS; 1999. Available from:
<https://pdfs.semanticscholar.org/104b/3127547393d6b94a8641100e9c297d653f56.pdf>

 3.
BAAR K. Training for Endurance and Strength. *Medicine & Science in Sports & Exercise*. 2006 Nov;38(11):1939–44.

 4.
Baar K, Hardie DG. Small molecules can have big effects on endurance. *Nature Chemical Biology*. 2008 Oct;4(10):583–4.

 5.
Barrès R, Yan J, Egan B, Treebak JT, Rasmussen M, Fritz T, et al. Acute Exercise Remodels Promoter Methylation in Human Skeletal Muscle. *Cell Metabolism*. 2012 Mar;15(3):405–11.

 6.
Carè A, Catalucci D, Felicetti F, Bonci D, Addario A, Gallo P, et al. MicroRNA-133 controls

cardiac hypertrophy. *Nature Medicine*. 2007 May;13(5):613–8.

7.

Chien KR. Molecular medicine: MicroRNAs and the tell-tale heart. *Nature*. 2007 May 24;447(7143):389–90.

8.

Eto Y, Yonekura K, Sonoda M, Arai N, Sata M, Sugiura S, et al. Calcineurin Is Activated in Rat Hearts With Physiological Left Ventricular Hypertrophy Induced by Voluntary Exercise Training. *Circulation*. 2000 May 9;101(18):2134–7.

9.

Fernandes T, Baraúna VG, Negrão CE, Phillips MI, Oliveira EM. Aerobic exercise training promotes physiological cardiac remodeling involving a set of microRNAs. *American Journal of Physiology-Heart and Circulatory Physiology*. 2015 Aug 15;309(4):H543–52.

10.

Iemitsu M, Maeda S, Jesmin S, Otsuki T, Kasuya Y, Miyauchi T. Activation pattern of MAPK signaling in the hearts of trained and untrained rats following a single bout of exercise. *Journal of Applied Physiology*. 2006 Jul;101(1):151–63.

11.

Maillet M, van Berlo JH, Molkenin JD. Molecular basis of physiological heart growth: fundamental concepts and new players. *Nature Reviews Molecular Cell Biology*. 2013 Jan;14(1):38–48.

12.

Wilkins BJ, Dai YS, Bueno OF, Parsons SA, Xu J, Plank DM, et al. Calcineurin/NFAT Coupling Participates in Pathological, but not Physiological, Cardiac Hypertrophy. *Circulation Research*. 2004 Jan 9;94(1):110–8.

13.

Boluyt MO, Brevick JL, Rogers DS, Randall MJ, Scalia AF, Li ZB. Changes in the rat heart proteome induced by exercise training: Increased abundance of heat shock protein hsp20. *PROTEOMICS*. 2006 May;6(10):3154-69.

14.

Burniston JG. Changes in the rat skeletal muscle proteome induced by moderate-intensity endurance exercise. *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics*. 2008 Jul;1784(7-8):1077-86.

15.

Burniston JG. Adaptation of the rat cardiac proteome in response to intensity-controlled endurance exercise. *PROTEOMICS*. 2009 Jan;9(1):106-15.

16.

Bye A, Langaas M, Høydal MA, Kemi OJ, Heinrich G, Koch LG, et al. Aerobic capacity-dependent differences in cardiac gene expression. *Physiological Genomics*. 2008 Mar;33(1):100-9.

17.

Bye A, Høydal MA, Catalucci D, Langaas M, Kemi OJ, Beisvag V, et al. Gene expression profiling of skeletal muscle in exercise-trained and sedentary rats with inborn high and low VO. *Physiological Genomics*. 2008 Nov;35(3):213-21.

18.

Iemitsu M, Maeda S, Miyauchi T, Matsuda M, Tanaka H. Gene expression profiling of exercise-induced cardiac hypertrophy in rats. *Acta Physiologica Scandinavica*. 2005 Nov 2;185(4):259-70.

19.

Kong SW, Bodyak N, Yue P, Liu Z, Brown J, Izumo S, et al. Genetic expression profiles during physiological and pathological cardiac hypertrophy and heart failure in rats. *Physiological Genomics*. 2005 Mar 21;21(1):34-42.

20.

Diffie GM. Adaptation of Cardiac Myocyte Contractile Properties to Exercise Training. *Exercise and Sport Sciences Reviews*. 2004 Jul;32(3):112-9.

21.

Kemi OJ, Haram PM, Wisløff U, Ellingsen Ø. Aerobic Fitness Is Associated With Cardiomyocyte Contractile Capacity and Endothelial Function in Exercise Training and Detraining. *Circulation*. 2004 Jun 15;109(23):2897-904.

22.

KEMI O, HARAM P, LOENNECHEN J, OSNES J, SKOMEDAL T, WISLOFF U, et al. Moderate vs. high exercise intensity: Differential effects on aerobic fitness, cardiomyocyte contractility, and endothelial function. *Cardiovascular Research*. 2005 Jul 1;67(1):161-72.

23.

Kemi OJ, Ellingsen Ø, Ceci M, Grimaldi S, Smith GL, Condorelli G, et al. Aerobic interval training enhances cardiomyocyte contractility and Ca²⁺ cycling by phosphorylation of CaMKII and Thr-17 of phospholamban. *Journal of Molecular and Cellular Cardiology*. 2007 Sep;43(3):354-61.

24.

Kemi OJ, Ceci M, Wisloff U, Grimaldi S, Gallo P, Smith GL, et al. Activation or inactivation of cardiac Akt/mTOR signaling diverges physiological from pathological hypertrophy. *Journal of Cellular Physiology*. 2008 Feb;214(2):316-21.

25.

Kemi OJ, Wisløff U. Mechanisms of exercise-induced improvements in the contractile apparatus of the mammalian myocardium. *Acta Physiologica*. 2010 Aug;199(4):425-39.

26.

Hsu CP, Huang CY, Wang JS, Sun PC, Shih CC. Extracellular Matrix Remodeling Attenuated After Experimental Postinfarct Left Ventricular Aneurysm Repair. *The Annals of Thoracic*

Surgery. 2008 Oct;86(4):1243-9.

27.

Burstein B, Nattel S. Atrial Fibrosis: Mechanisms and Clinical Relevance in Atrial Fibrillation. *Journal of the American College of Cardiology*. 2008 Feb;51(8):802-9.

28.

KOVANEN V, SUOMINEN H, HEIKKINEN E. Connective tissue of "fast" and "slow" skeletal muscle in rats...effects of endurance training. *Acta Physiologica Scandinavica*. 1980 Feb;108(2):173-80.

29.

Daniels A, van Bilsen M, Goldschmeding R, van der Vusse GJ, van Nieuwenhoven FA. Connective tissue growth factor and cardiac fibrosis. *Acta Physiologica*. 2009 Mar;195(3):321-38.

30.

Creemers EEJM, Davis JN, Parkhurst AM, Leenders P, Dowdy KB, Hapke E, et al. Deficiency of TIMP-1 exacerbates LV remodeling after myocardial infarction in mice. *American Journal of Physiology-Heart and Circulatory Physiology*. 2003 Jan;284(1):H364-71.

31.

Williams PE, Goldspink G. Connective tissue changes in immobilised muscle. 138(2):343-50. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1164074/>

32.

MURPHY G, NAGASE H. Progress in matrix metalloproteinase research. *Molecular Aspects of Medicine*. 2008 Oct;29(5):290-308.

33.

Di Biase V, Franzini-Armstrong C. Evolution of skeletal type e-c coupling. *The Journal of Cell Biology*. 2005 Nov 21;171(4):695-704.

34.

Meeusen R, Piacentini MF, Busschaert B, Buyse L, De Schutter G, Stray-Gundersen J. Hormonal responses in athletes: the use of a two bout exercise protocol to detect subtle differences in (over)training status. *European Journal of Applied Physiology*. 2004 Mar;91(2-3):140-6.

35.

BOOTH FW, TSENG BS, FLUCK M, CARSON JA. Molecular and cellular adaptation of muscle in response to physical training. *Acta Physiologica Scandinavica*. 1998 Feb;162(3):343-50.

36.

Hill M, Wernig A, Goldspink G. Muscle satellite (stem) cell activation during local tissue injury and repair. *Journal of Anatomy*. 2003 Jul;203(1):89-99.

37.

Reid MB. Response of the ubiquitin-proteasome pathway to changes in muscle activity. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 2005 Jun;288(6):R1423-31.

38.

Hambrecht R, Adams V, Erbs S, Linke A,
Kra

..
nkel N, Shu Y, et al. Regular Physical Activity Improves Endothelial Function in Patients With Coronary Artery Disease by Increasing Phosphorylation of Endothelial Nitric Oxide Synthase. *Circulation*. 2003 Jul;107(25):3152-8.

39.

Haram PM, Adams V, Kemi OJ, Brubakk AO, Hambrecht R, Ellingsen Ø, et al. Time-course of endothelial adaptation following acute and regular exercise. *European Journal of*

Cardiovascular Prevention & Rehabilitation. 2006 Aug;13(4):585–91.

40.

Haram PM, Kemi OJ, Wisloff U. Adaptation of endothelium to exercise training: Insights from experimental studies. *1AD*;13:336–46. Available from: <https://www.bioscience.org/2008/v13/af/2683/fulltext.htm>

41.

Linke A, Erbs S, Hambrecht R. Effects of exercise training upon endothelial function in patients with cardiovascular disease. *1AD*;13:424–32. Available from: <https://www.bioscience.org/2008/v13/af/2689/fulltext.htm>

42.

Miyachi M, Iemitsu M, Okutsu M, Onodera S. Effects of endurance training on the size and blood flow of the arterial conductance vessels in humans. *Acta Physiologica Scandinavica*. 1998 May;163(1):13–6.

43.

Spence AL, Carter HH, Naylor LH, Green DJ. A prospective randomized longitudinal study involving 6 months of endurance or resistance exercise. Conduit artery adaptation in humans. *The Journal of Physiology*. 2013 Mar;591(5):1265–75.

44.

Bogdanis GC, Nevill ME, Boobis LH, Lakomy HK, Nevill AM. Recovery of power output and muscle metabolites following 30 s of maximal sprint cycling in man. *The Journal of Physiology*. 1995 Jan 15;482(2):467–80.

45.

Casey A, Constantin-Teodosiu D, Howell S, Hultman E, Greenhaff PL. Creatine ingestion favorably affects performance and muscle metabolism during maximal exercise in humans. *American Journal of Physiology-Endocrinology and Metabolism*. 1996 Jul;271(1):E31–7.

46.

Jørgensen SB, Richter EA, Wojtaszewski JFP. Role of AMPK in skeletal muscle metabolic regulation and adaptation in relation to exercise. *The Journal of Physiology*. 2006 Jul 1;574(1):17–31.

47.

Kiens B, Richter EA. Utilization of skeletal muscle triacylglycerol during postexercise recovery in humans. *American Journal of Physiology-Endocrinology and Metabolism*. 1998 Aug;275(2):E332–7.

48.

Tsintzas OK, Williams C, Boobis L, Greenhaff P. Carbohydrate ingestion and single muscle fiber glycogen metabolism during prolonged running in men. *Journal of Applied Physiology*. 1996 Aug;81(2):801–9.

49.

Walter G, Vandeborne K, McCully KK, Leigh JS. Noninvasive measurement of phosphocreatine recovery kinetics in single human muscles. *American Journal of Physiology-Cell Physiology*. 1997 Feb;272(2):C525–34.

50.

KEMI O, HOYDAL M, HARAM P, GARNIER A, FORTIN D, VENTURACLAPIER R, et al. Exercise training restores aerobic capacity and energy transfer systems in heart failure treated with losartan. *Cardiovascular Research*. 2007 Oct 1;76(1):91–9.

51.

Wisløff U. Aerobic exercise reduces cardiomyocyte hypertrophy and increases contractility, Ca²⁺ sensitivity and SERCA-2 in rat after myocardial infarction. *Cardiovascular Research*. 2002 Apr;54(1):162–74.

52.

Wisløff U, Støylen A, Loennechen JP, Bruvold M, Rognum Ø, Haram PM, et al. Superior Cardiovascular Effect of Aerobic Interval Training Versus Moderate Continuous Training in

Heart Failure Patients. *Circulation*. 2007 Jun 19;115(24):3086–94.

53.

Hawley JA, Hargreaves M, Joyner MJ, Zierath JR. Integrative Biology of Exercise. *Cell*. 2014 Nov;159(4):738–49.

54.

Rowe GC, Safdar A, Arany Z. Running Forward. *Circulation*. 2014 Feb 18;129(7):798–810.